

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 October 2003 (09.10.2003)

PCT

(10) International Publication Number
WO 03/082291 A1

- (51) International Patent Classification⁷: **A61K 31/5375**,
C07D 265/30
- (21) International Application Number: PCT/EP03/03339
- (22) International Filing Date: 27 March 2003 (27.03.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0207449.0 28 March 2002 (28.03.2002) GB
- (71) Applicant (for all designated States except US): **GLAXO GROUP LIMITED** [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **ANCLIFF, Rachael, Ann** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **COOK, Caroline, Mary** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **ELDRED, Colin, David** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **GORE, Paul, Martin** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **HARRISON, Lee, Andrew** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **HAYES, Martin, Alistair** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **HODGSON, Simon,**


Teanby [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **JUDD, Duncan, Bruce** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **KEELING, Suzanne, Elaine** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **LEWELL, Xiao, Qing** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **MILLS, Gail** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **ROBERTSON, Graeme, Michael** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **SWANSON, Stephen** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **WALKER, Andrew, John** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **WILKINSON, Mark** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB).

(74) Agent: **GIDDINGS, Peter, John**; GlaxoSmithKline, Corporate Intellectual Property (CN925.1), 980 Great West Road, Brentford, Middlesex TW8 9GS (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

[Continued on next page]

(54) Title: N-{ (2S)-4-(3,4-DIFLUOROBENZYL)MORPHOLIN-2YL}METHYL-2-{3-(METHYLSULPHONYL)AMINO}PHENYL}ACETAMIDE AS CCR3 ANTAGONIST FOR THE TREATMENT OF INFLAMMATORY CONDITIONS



(57) Abstract: Compound of formula (I): HsC H H '0) IIZZ V which is; 3-(f (((2S)-4-(3,4-dichlorobenzyl)morpholin-2-yl)methyl)amino)carbonyl)-amino)methyl)=Nethylbenzamide; and salts and solvates thereof, a process for its preparation intermediates, pharmaceutical formulations containing it and its use in therapy, as CCR3=antagonist for the treatment of inflammatory conditions.



(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

N-{4-(3,4-DIFLUOROBENZYL)MORPHOLIN-2-YLMETHYL}-2-{3-(METHYLSULPHONYLAMINO)PHENYL}ACETAMIDE AS CCR3 ANTAGONIST FOR THE TREATMENT OF INFLAMMATORY CONDITIONS

Novel Compounds

This invention relates to a novel compound, a process for its preparation, pharmaceutical formulations containing it and its use in therapy.

5 Inflammation is a primary response to tissue injury or microbial invasion and is characterised by leukocyte adhesion to the endothelium, diapedesis and activation within the tissue. Leukocyte activation can result in the generation of toxic oxygen species (such as superoxide anion), and the release of granule products (such as peroxidases and proteases). Circulating leukocytes include
 10 neutrophils, eosinophils, basophils, monocytes and lymphocytes. Different forms of inflammation involve different types of infiltrating leukocytes, the particular profile being regulated by the profile of adhesion molecule, cytokine and chemotactic factor expression within the tissue.

The primary function of leukocytes is to defend the host from invading
 15 organisms, such as bacteria and parasites. Once a tissue is injured or infected, a series of events occurs which causes the local recruitment of leukocytes from the circulation into the affected tissue. Leukocyte recruitment is controlled to allow for the orderly destruction and phagocytosis of foreign or dead cells, followed by tissue repair and resolution of the inflammatory infiltrate. However in
 20 chronic inflammatory states, recruitment is often inappropriate, resolution is not adequately controlled and the inflammatory reaction causes tissue destruction.

There is increasing evidence that the bronchial inflammation which is characteristic of asthma represents a specialised form of cell-mediated immunity, in which cytokine products, such as IL-4 and IL-5 released by T-helper 2 (Th2)
 25 lymphocytes, orchestrate the accumulation and activation of granulocytes, in particular eosinophils and to a lesser extent basophils. Through the release of cytotoxic basic proteins, pro-inflammatory mediators and oxygen radicals, eosinophils generate mucosal damage and initiate mechanisms that underlie bronchial hyperreactivity. Therefore, blocking the recruitment and activation of
 30 Th2 cells and eosinophils is likely to have anti-inflammatory properties in asthma. In addition, eosinophils have been implicated in other disease types such as rhinitis, eczema, irritable bowel syndrome and parasitic infections.

Chemokines are a large family of small proteins which are involved in trafficking and recruitment of leukocytes (for review see Luster, New Eng. J.
 35 Med., 338, 436-445 (1998)). They are released by a wide variety of cells and act to attract and activate various cell types, including eosinophils, basophils, neutrophils, macrophages, T and B lymphocytes. There are two major families of chemokines, CXC- (α) and CC- (β) chemokines, classified according to the spacing of two conserved cysteine residues near to the amino terminus of the
 40 chemokine proteins. Chemokines bind to specific cell surface receptors

PG4791

belonging to the family of G-protein-coupled seven transmembrane-domain proteins (for review see Luster, 1998). Activation of chemokine receptors results in, amongst other responses, an increase in intracellular calcium, changes in cell shape, increased expression of cellular adhesion molecules, degranulation and
5 promotion of cell migration (chemotaxis).

To date a number of CC chemokine receptors have been identified and of particular importance to the current invention is the CC-chemokine receptor-3 (CCR-3), which is predominantly expressed on eosinophils, and also on basophils, mast cells and Th2 cells. Chemokines that act at CCR-3, such as
10 RANTES, MCP-3 and MCP-4, are known to recruit and activate eosinophils. Of particular interest are eotaxin and eotaxin-2, which specifically bind to CCR-3. The localization and function of CCR-3 chemokines indicate that they play a central role in the development of allergic diseases such as asthma. Thus, CCR-3 is specifically expressed on all the major cell types involved in inflammatory
15 allergic responses. Chemokines that act at CCR-3 are generated in response to inflammatory stimuli and act to recruit these cell types to sites of inflammation, where they cause their activation (e.g. Griffiths et al., J. Exp. Med., 179, 881-887 (1994), Lloyd et al., J. Exp. Med., 191, 265-273 (2000)). In addition, anti-CCR-3 monoclonal antibodies completely inhibit eotaxin interaction with eosinophils
20 (Heath, H. *et al.*, J. Clin. Invest. 99 (2), 178-184 (1997)), while an antibody for the CCR-3 specific chemokine, eotaxin, reduced both bronchial hyperreactivity and lung eosinophilia in an animal model of asthma (Gonzalo et al., J. Exp. Med., 188, 157-167 (1998). Thus, many lines of evidence indicate that antagonists at the CCR-3 receptor are very likely to be of therapeutic use for the treatment of a
25 range of inflammatory conditions.


In addition to a key role in inflammatory disorders, chemokines and their receptors also play a role in infectious disease. Mammalian cytomegaloviruses, herpes viruses and pox viruses express chemokine receptor homologues, which can be activated by human CC chemokines such as RANTES and MCP-3
30 receptors (for review see Wells and Schwartz, Curr. Opin. Biotech., 8, 741-748, 1997). In addition, human chemokine receptors, such as CXCR-4, CCR-5 and CCR-3, can act as co-receptors for the infection of mammalian cells by microbes such as human immunodeficiency viruses (HIV). Thus, chemokine receptor antagonists, including CCR-3 antagonists, may be useful in blocking infection of
35 CCR-3 expressing cells by HIV or in preventing the manipulation of immune cellular responses by viruses such as cytomegaloviruses.

International Patent Application publication number WO 01/24786 (Shionogi & Co. Ltd.) discloses certain aryl and heteroaryl derivatives for treating diabetes. WO 00/69830 (Torrey Pines Institute for Molecular Studies) discloses
40 certain diazacyclic compounds, and libraries containing them, for biological

PG4791

- screening. WO 00/18767 (Neurogen Corporation) discloses certain piperazine derivatives as dopamine D4 receptor antagonists. United States Patent 6,031,097 and WO 99/21848 (Neurogen Corporation) discloses certain aminoisoquinoline derivatives as dopamine receptor ligands. WO 99/06384
- 5 (Recordati Industria Chimica) discloses piperazine derivatives useful for the treatment of neuromuscular dysfunction of the lower urinary tract. WO 98/56771 (Schering Aktiengesellschaft) discloses certain piperazine derivatives as anti-inflammatory agents. WO 97/47601 (Yoshitomi Pharmaceutical Industries Ltd.) discloses certain fused heterocyclic compounds as dopamine D-receptor
- 10 blocking agents. WO 96/39386 (Schering Corporation) discloses certain piperidine derivatives as neurokinin antagonists. WO 96/02534 (Byk Gulden Lomberg Chemische Fabrik GmbH) discloses certain piperazine thiopyridines useful for controlling helicobacter bacteria. WO 95/32196 (Merck Sharp & Dohme Limited) discloses certain piperazine, piperidine, and tetrahydropyridine
- 15 derivatives as 5-HT1D-alpha antagonists. United States Patent 5,389,635 (E.I. Du Pont de Nemours and Company) discloses certain substituted imadazoles as angiotensin-II antagonists. European Patent Application publication number 0 306 440 (Schering Aktiengesellschaft) discloses certain imidazole derivatives as cardiovascular agents.
- 20 A novel compound has now been found which is a CCR-3 antagonist. This compound blocks the migration/chemotaxis of eosinophils and thus possesses anti-inflammatory properties. This compound is therefore of potential therapeutic benefit, especially in providing protection from eosinophil, basophil and Th2-cell-induced tissue damage in diseases where such cell types are
- 25 implicated, particularly allergic diseases, including but not limited to bronchial asthma, allergic rhinitis and atopic dermatitis.

Thus, according to one aspect of the invention, there is provided a compound of formula (I):



which is;

PG4791


N-[[[(2S)-4-(3,4-difluorobenzyl)morpholin-2-yl]methyl]-2-{3-
[(methylsulfonyl)amino]phenyl}acetamide;
and salts and solvates thereof.

Suitable salts of the compound of formula (I) include physiologically
5 acceptable salts and salts which may not be physiologically acceptable but may
be useful in the preparation of compounds of formula (I) and physiologically
acceptable salts thereof. If appropriate, acid addition salts may be derived from
inorganic or organic acids, for example hydrochlorides, hydrobromides,
sulphates, phosphates, acetates, benzoates, citrates, succinates, lactates,
10 tartrates, fumarates, maleates, 1-hydroxy-2-naphthoates, pamoates,
methanesulphonates, formates or trifluoroacetates.

Examples of solvates include hydrates.


The compound of formula (I) and salts and solvates thereof may be
prepared by the methodology described hereinafter, constituting a further aspect
15 of this invention.

A process according to the invention for the preparation of the compound
of formula (I) comprises coupling of a compound of formula (II) or a salt thereof:



20

with a compound of formula (III):



25 and thereafter, if required, carrying out one or more of the following optional
steps:

- (i) removing any protecting group(s);
- (ii) preparing an appropriate salt or solvate of the compound so formed.


PG4791

The coupling of the compounds of formulae (II) and (III) is carried out in any suitable solvent, for example a polar organic solvent such as N,N-dimethylformamide in the presence of a suitable dehydrating agent such as a carbodiimide reagent e.g. 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, and optionally a suitable activating agent, e.g. an activated
 5 hydroxy compound such as 1-hydroxybenzotriazole, and optionally in the presence of a base such as a tertiary amine, e.g. N,N-diisopropylethylamine, under conventional coupling conditions at any temperature providing a suitable rate of formation of the required product, over a suitable reaction time, for
 10 example 12 - 24 hours.

Suitable reaction temperatures include those in the range of 18°C to 25°C.

The reaction products are isolated and purified using conventional methods.


15 The compound of formula (III) may be prepared by sulphonylation of the compound of formula (IV):



20 The sulphonylation may be carried out using a conventional sulphonylating agent, for example a sulphonyl chloride such as methanesulphonyl chloride, in the presence of a suitable base, for example aqueous sodium carbonate.


The compound of formula (II) may be prepared by either by Reaction (a),
 25 Reaction (b), or Reaction (c).

Reaction (a). Reaction of the compound of formula (V) with a compound of formula (VI)



30 wherein A is a protected amino group, suitably phthalimido, followed by deprotection of the amino group to give a compound of formula (IIR)

PG4791



followed by resolution of the resulting enantiomers of the compound of formula (IIR);

5 or;


Reaction (b). Reaction of a compound of formula (V) as hereinbefore defined with a compound of formula (VIA)



10

wherein A is as hereinbefore defined for formula (VI), followed by deprotection of the amino group to give the compound of formula (II).

Reaction (c). Hydrolysis of the compound of formula (VII);



15

followed by resolution of the resulting enantiomers of a compound of formula (IIR).

For both reactions (a) and (b), the cyclisation of the intermediate diols (IIBR) and (IIB) in the reaction between the compound of formula (V) and a compound of formula (VI) or (VIA) is typically carried out under the Mitsunobu conditions as follows:

Typically, a mixture of the compound of formula (V) and the compound of formula (VI) or formula (VIA) in a suitable solvent, such as tetrahydrofuran, is stirred, suitably for 20-24 hours at a suitable temperature, suitably the reflux temperature of the solvent, under an inert atmosphere, suitably an atmosphere of



PG4791

nitrogen. Further solvent is then added and the mixture cooled, suitably to 0-5°C. A suitable phosphine, suitably triphenyl phosphine, is added and the mixture stirred until all the solid is dissolved. A suitable azo compound, suitably diisopropylazodicarboxylate, is then added over a period of time, suitably, 10-15 minutes, while maintaining the temperature at <7°C. The mixture is allowed to stand for a period of time, suitably 2-3 hours, then allowed to warm, suitably to 20-25°C. After a further period of standing, suitably 4-6 hours, further phosphine and azo compound are added. After a further period of standing, suitably 20-24 hours, the reaction mixture is concentrated to near dryness. A suitable alcohol, suitably propan-2-ol, is added and the concentration step repeated; the alcohol addition and concentration step is then repeated. Further alcohol is then added and the mixture heated to a temperature suitably between 65-75°C. After a suitable period, suitably 20-45 minutes, the resultant slurry is cooled suitably to 20-25°C, and then allowed to stand, suitably for 1.5 – 3 hours, after which time the product is isolated by filtration. The filter bed is washed with more alcohol and then dried *in vacuo* at 35-45°C to yield the protected form of the compound of formula (IIR) or formula (II) respectively.

The removal of the protecting group from the product is typically carried out as follows. A slurry of the protected form of the compound of formula (IIR) or formula (II) in an appropriate polar solvent, suitably water, is heated to elevated temperature, suitably 70-75°C and then treated dropwise with a concentrated mineral acid, suitably concentrated sulphuric acid. The mixture was then heated at elevated temperature, suitably the reflux temperature of the solvent, for a suitable period of time, suitably 20-24 hours, after which the reaction mixture was cooled to 20-25°C and then treated with a suitable apolar solvent, suitably dichloromethane. A base, suitably 0.880 ammonia solution, is then added dropwise, maintaining the temperature between 20-25°C. Further apolar solvent is then added, the aqueous phase then being separated and extracted with further apolar solvent. The combined organic phase is washed with water and then evaporated to dryness. The residue is redissolved and the apolar solvent evaporated to give the compound of formula (IIR) or formula (II).

The process for the preparation of the protected form of the compound of formula (IIR) or formula (II) described above may also be undertaken in two stages, in which an intermediate compound of formula (IIBR) or of formula (IIB) respectively;

PG4791



wherein A is as hereinbefore defined for formulae (VI) and (VIA);
is isolated.

- 5 Typically, a mixture of the compound of formula (V) and a compound of formula (VI) or formula (VIA) in a suitable solvent, such as tetrahydrofuran, is stirred, suitably for 20-24 hours at a suitable temperature, suitably the reflux temperature of the solvent, under an inert atmosphere, suitably an atmosphere of nitrogen. Further compound of formula (V) is added and the mixture heated at a
- 10 suitable temperature, suitably the reflux temperature of the solvent, under an inert atmosphere, suitably an atmosphere of nitrogen, for a suitable period of time, suitably 3-6 hours. The reaction mixture is then cooled, suitably to 20-25°C, and the compound precipitated by means of addition of a suitable co-solvent, suitably diisopropyl ether. The compound of formula (IIBR) or formula
- 15 (IIB) respectively is isolated by filtration, washed with further co-solvent and dried *in vacuo*.

- A protected form of the compound of formula (IIR) or formula (II) may then be prepared from a compound of formula (IIBR) or formula (IIB) under similar conditions to those of the reaction between a compound of formula (V)
- 20 and formulae (VI) or (VIA) as hereinbefore described, but omitting the reflux period prior to the addition of the phosphine and azo compounds.


- Reaction (c) is typically carried out by stirring a solution of the compound of formula (VII) in a suitable solvent, for example a mixture of methanol and water, and adding a suitable base, for example potassium carbonate. The
- 25 mixture is stirred at a suitable temperature, for example those in the range 20-25°C for a suitable time, for example 16-20 hours followed by removal of the organic solvent *in vacuo*. Water is then added and the mixture extracted with a suitable organic solvent, for example ethyl acetate. The combined organic phases are washed with water and saturated aqueous sodium chloride solution
- 30 before drying over a suitable drying agent, for example sodium sulphate, filtering and evaporation of the solvent *in vacuo*. The crude product is then purified by flash chromatography.

The resolution of the compound of formula (II) from the racemic product i.e. the compound of formula (IIR) may be undertaken using techniques well

PG4791


known to those skilled in the art, for example preparative chiral high performance liquid chromatography (chiral HPLC) or by fractional crystallisation of diastereoisomeric salts.

The compound of formula (V) may be prepared from the compound of
5 formula (IX)



by reaction with ethanolamine. Suitably the reaction is carried out at elevated
10 temperature, e.g. 40 - 60°C, in the absence of solvent.

The compound of formula (VII) may be prepared by reaction of the compound of formula (VIII)



15 with 3,4-difluorobenzyl chloride.

The reaction between the compound of formula (VIII) and 3,4-difluorobenzyl chloride is typically carried out in a suitable solvent, for example N,N-dimethylformamide, under an inert atmosphere, for example an atmosphere
20 of nitrogen, with the addition of a suitable base, for example potassium carbonate, and a suitable activating agent, such as sodium iodide. The mixture is then stirred at a suitable temperature, for example a temperature in the range of 20-25°C, for a suitable period of time, for example 16-20 hours before removing the volatile components in vacuo.

25 The compound of formula (VII) is prepared by treating a solution of morpholin-2-ylmethylamine in a suitable organic solvent, for example methanol, under an inert atmosphere, for example an atmosphere nitrogen, with a solution of ethyl- α,α,α -trifluoroacetate in a suitable organic solvent, for example ether. The mixture is then stirred for a suitable period of time, for example 20-40
30 minutes at a suitable temperature, for example a temperature in the range of 20-25°C and the volatile components removed in vacuo. The residue is then dissolved in a suitable organic solvent, for example methanol, and the volatile components removed in vacuo.

PG4791

The compounds of formula (IV) and (VI) are commercially available compounds and may be prepared by analogy with known procedures, for example those disclosed in standard reference texts of synthetic methodology such as *J. March, Advanced Organic Chemistry, 3rd Edition (1985), Wiley*

5 *Interscience.*

The compounds of formulae (II), (IIBR), (IIB) and (V) are considered to be novel.

Accordingly, there is provided a compound of formula (II).

There is also provided a compound of formula (IIBR).

10 There is also provided a compound of formula (IIB).

There is also provided a compound of formula (V).

Suitable protecting groups in any of the above mentioned reactions are those used conventionally in the art. The methods of formation and removal of such protecting groups are those conventional methods appropriate to the molecule being protected, for example those methods discussed in standard reference texts of synthetic methodology such as *P J Kocienski, Protecting Groups, (1994), Thieme.*

For any of the hereinbefore described reactions or processes, conventional methods of heating and cooling may be employed, for example electric hotplates and ice/salt baths respectively. Conventional methods of purification, for example crystallisation and column chromatography may be used as required.

The salts and solvates of the compounds of formula (I) may be prepared from the compound of formula (I) or suitable salts or solvates thereof and isolated according to conventional procedures.

The compounds of the invention may be tested for *in vitro* biological activity in accordance with the following assays:

(a) CCR-3 Binding Assay

30 A CCR-3 competition binding SPA (scintillation proximity assay) was used to assess the affinity of novel compounds for CCR-3. Membranes prepared from K562 cells stably expressing CCR-3 (2.5µg/well) were mixed with 0.25mg/well wheat-germ agglutinin SPA beads (Amersham) and incubated in binding buffer (HEPES 50 mM, CaCl₂ 1 mM, MgCl₂ 5 mM, 0.5% BSA) at 4°C for 1.5 hr. Following incubation, 20 pM of [¹²⁵I] eotaxin (Amersham) and increasing concentrations of compound (1pM to 30µM) were added and incubated in a 96 well plate for 2 hr at 22°C then counted on a Microbeta plate counter. The total assay volume was 100 µl. Competition binding data were analysed by fitting the data with a four parameter logistic equation. Data are presented as the mean

PG4791

pIC₅₀ values (negative logarithm of the concentration of compound which inhibits [¹²⁵I]eotaxin binding by 50%) from at least two experiments.

(b) Eosinophil Chemotaxis Assay.

- 5 The compound was evaluated for its inhibitory effect on eosinophil chemotaxis. Eosinophils were purified from human peripheral blood by standard CD16 cell depletion using a Miltenyi cell separation column and a magnetic Super Macs magnet as previously described (Motegi & Kita, 1998; J.Immunology. 161:4340-6). Cells were re-suspended in RPMI 1640/10% FCS
- 10 solution and incubated with calcein-AM (Molecular Probes) at 37°C for 30 mins. Following incubation, the eosinophils were centrifuged at 400g for 5 min and re-suspended in RPMI/FCS at 2.2 million/ml. Cells were then incubated in the presence of increasing concentration of compounds (1 pM to 30 µM) at 37°C for 30 mins. For control responses cells were incubated with RPMI/FCS only. The
- 15 agonist eotaxin (an EC₈₀ concentration) was added to the lower chamber of a 96 well chemotaxis plate (5 µm filter: Receptor Technologies). Eosinophils (50 µl of 2 million/ml cells) were added to the top chamber of the filter plate and incubated at 37°C for 45 mins. Cells remaining on top of the chemotaxis filter were removed and the number of eosinophils which had migrated were quantified by
- 20 reading the plate on a fluorescent plate reader. Inhibition curves for the effect of compounds on eosinophil chemotaxis were analysed by fitting the data with a four parameter logistic equation. Functional pK_i values (fpK_i) were generated using the equation below (Lazareno & Birdsall, 1995. Br.J.Pharmacol 109: 1110-9).

25

$$fpKi = \frac{IC_{50}}{1 + \frac{[Agonist]}{EC_{50}}}$$

- The compound of the invention was tested in the CCR-3 binding and/or eosinophil chemotaxis assays (assays (a) and (b)). The compound of the
- 30 invention was tested in the CCR-3 binding assay and possessed a pIC₅₀ value of 8.0. The compound of the invention tested in the CCR-3 eosinophil chemotaxis assay possessed an fpK_i value of 8.4.

- Examples of disease states in which the compound of the invention has potentially beneficial anti-inflammatory effects include diseases of the respiratory
- 35 tract such as bronchitis (including chronic bronchitis), bronchiectasis, asthma (including allergen-induced asthmatic reactions), chronic obstructive pulmonary disease (COPD), cystic fibrosis, sinusitis and rhinitis.

PG4791

Also included are diseases of the gastrointestinal tract such as intestinal inflammatory diseases including inflammatory bowel disease (e.g. Crohn's disease or ulcerative colitis) and intestinal inflammatory diseases secondary to radiation exposure or allergen exposure.

5 Furthermore, the compound of the invention may be used to treat nephritis; skin diseases such as psoriasis, eczema, allergic dermatitis and hypersensitivity reactions; and diseases of the central nervous system which have an inflammatory component (eg. Alzheimer's disease, meningitis, multiple sclerosis), HIV and AIDS dementia.

10 The compounds of the present invention may also be of use in the treatment of nasal polyposis, conjunctivitis or pruritis.

Further examples of disease states in which the compound of the invention has potentially beneficial effects include cardiovascular conditions such as atherosclerosis, peripheral vascular disease and idiopathic hypereosinophilic
15 syndrome.

The compounds of the invention may be useful as an immunosuppressive agents and so has use in the treatment of auto-immune diseases such as allograft tissue rejection after transplantation, rheumatoid arthritis and diabetes. The compound of the invention may also be useful in inhibiting metastasis.

20 Diseases of principal interest include asthma, COPD and inflammatory diseases of the upper respiratory tract involving seasonal and perennial rhinitis.

It will be appreciated by those skilled in the art that reference herein to treatment extends to prophylaxis as well as the treatment of established conditions.

25 As mentioned above, the compound of formula (I) is useful as a therapeutic agent.

There is thus provided as a further aspect of the invention a compound of formula (I) or a physiologically acceptable salt or solvate thereof for use as an active therapeutic agent.

30 There is also therefore provided the compound of formula (I), or a physiologically acceptable salt or solvate thereof, for use in the treatment of inflammatory conditions, eg. asthma or rhinitis.

According to another aspect of the invention, there is provided the use of the compound of formula (I) or a physiologically acceptable salt or solvate
35 thereof for the manufacture of a medicament for the treatment of patients with inflammatory conditions, eg. asthma or rhinitis.

In a further or alternative aspect there is provided a method for the treatment of a human or animal subject with an inflammatory condition eg. asthma or rhinitis, which method comprises administering an effective amount of

PG4791

the compound of formula (I) or a physiologically acceptable salt or solvate thereof.

The compound according to the invention may be formulated for administration in any convenient way.

- 5 There is thus further provided a pharmaceutical composition comprising the compound of formula (I), or a physiologically acceptable salt or solvate thereof, and optionally one or more physiologically acceptable diluents or carriers.

- There is also provided a process for preparing such a pharmaceutical formulation which comprises admixing the compound of formula (I) or a
10 physiologically acceptable salt or solvate thereof with one or more physiologically acceptable diluents or carriers.

The compound according to the invention may, for example, be formulated for oral, inhaled, intranasal, buccal, parenteral or rectal administration, preferably for oral administration.

- 15 Tablets and capsules for oral administration may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, mucilage of starch, cellulose or polyvinyl pyrrolidone; fillers, for example, lactose, microcrystalline cellulose, sugar, maize- starch, calcium phosphate or sorbitol; lubricants, for example, magnesium stearate, stearic acid,
20 talc, polyethylene glycol or silica; disintegrants, for example, potato starch, croscarmellose sodium or sodium starch glycollate; or wetting agents such as sodium lauryl sulphate. The tablets may be coated according to methods well known in the art.

- Oral liquid preparations may be in the form of, for example, aqueous or
25 oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example, sorbitol syrup, methyl cellulose, glucose/sugar syrup, gelatin, hydroxymethyl cellulose, carboxymethyl cellulose, aluminium stearate gel
30 or hydrogenated edible fats; emulsifying agents, for example, lecithin, sorbitan mono-oleate or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, fractionated coconut oil, oily esters, propylene glycol or ethyl alcohol; or preservatives, for example, methyl or propyl *p*- hydroxybenzoates or sorbic acid. The preparations may also contain buffer salts, flavouring, colouring
35 and/or sweetening agents (e.g. mannitol) as appropriate.

For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

The compound may also be formulated as suppositories, e.g. containing conventional suppository bases such as cocoa butter or other glycerides.

PG4791

The compound according to the invention may also be formulated for parenteral administration by bolus injection or continuous infusion and may be presented in unit dose form, for instance as ampoules, vials, small volume infusions or pre-filled syringes, or in multidose containers with an added
5 preservative. The compositions may take such forms as solutions, suspensions, or emulsions in aqueous or non-aqueous vehicles, and may contain formulatory agents such as anti-oxidants, buffers, antimicrobial agents and/or tonicity adjusting agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile, pyrogen-free water, before use.
10 The dry solid presentation may be prepared by filling a sterile powder aseptically into individual sterile containers or by filling a sterile solution aseptically into each container and freeze-drying.

The compound and pharmaceutical compositions according to the invention may also be used in combination with other therapeutic agents, for
15 example antihistaminic agents, anticholinergic agents, anti-inflammatory agents such as corticosteroids, e.g. fluticasone propionate, beclomethasone dipropionate, mometasone furoate, triamcinolone acetonide or budesonide; or non-steroidal anti-inflammatory drugs (NSAIDs) eg. sodium cromoglycate, nedocromil sodium, PDE-4 inhibitors, leukotriene antagonists, iNOS inhibitors,
20 tryptase and elastase inhibitors, beta-2 integrin antagonists and adenosine 2a agonists; or beta adrenergic agents such as salmeterol, salbutamol, formoterol, fenoterol or terbutaline and salts thereof; or antiinfective agents e.g. antibiotic agents and antiviral agents. It will be appreciated that when the compound of the present invention is administered in combination with other therapeutic agents
25 normally administered by the inhaled or intranasal route, that the resultant pharmaceutical composition may be administered by the inhaled or intranasal route.

The compounds of the invention may conveniently be administered in amounts of, for example, 0.001 to 500mg/kg body weight, preferably 0.01 to
30 500mg/kg body weight, more preferably 0.01 to 100mg/kg body weight, and at any appropriate frequency e.g. 1 to 4 times daily. The precise dosing regimen will of course depend on factors such as the therapeutic indication, the age and condition of the patient, and the particular route of administration chosen.

Throughout the description and the claims which follow, unless the
35 context requires otherwise, the word 'comprise', and variations such as 'comprises' and 'comprising', will be understood to imply the inclusion of a stated integer or step or group of integers but not to the exclusion of any other integer or step or group of integers or steps.

40 General Experimental Details

PG4791

Liquid Chromatography Mass Spectrometry (LC/MS) System

The following (LC/MS) system was used:

an 3µm ABZ+PLUS (3.3cm x 4.6mm internal diameter) column, eluting with solvents: A – 0.1%v/v formic acid + 0.077% w/v ammonium acetate in water; and

- 5 B – 95:5 acetonitrile:water + 0.05%v/v formic acid, at a flow rate of 3 ml per minute. The following gradient protocol was used: 100% A for 0.7mins; A+B mixtures, gradient profile 0 – 100% B over 3.5mins; hold at 100%B for 1.1mins; return to 100% A over 0.2mins.


Solid phase extraction (ion exchange)

- 10 'SCX' refers to Isolute Flash SCX-2 sulphonic acid solid phase extraction cartridges.

All temperatures are in °C

Example 1: N-[(2S)-4-(3,4-difluorobenzyl)morpholin-2-yl]methyl]-2-{3-


- 15 [(methylsulfonyl)amino]phenyl}acetamide



- To a stirred solution of Description 4 (0.0242g) in N,N-dimethylformamide (1ml)
 20 was added a solution of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.0304g), 1-hydroxybenzotriazole (0.0171g), and Description 3 (0.0243g) in N,N-dimethylformamide (1ml) at 22°. N,N-diisopropylethylamine (0.0368ml) was added to the mixture which was then stirred at 22° for 18h. The mixture was applied to a 2g SCX ion-exchange cartridge (IST Isolute Flash SCX-
 25 2), pre-conditioned with methanol. The cartridge was eluted with methanol and 10% 0.880 ammonia solution in methanol. The first ammonia fraction was evaporated *in vacuo* and the residue was further purified by Biotage™ flash chromatography on silica gel, eluting with 200:8:1 dichloromethane/ethanol/0.880 ammonia solution. The required fractions were combined and the solvent
 30 evaporated *in vacuo* to give the title compound (0.0353g) as a colourless glass.
 LC/MS : R_t = 2.16min, m/z 454 [MH⁺]


Description 1: 2-(3,4-Difluorobenzylamino)-ethanol

PG4791



- 4-Chloromethyl-1,2-difluorobenzene (90.0g) (Fluorochem chemicals/Journal of Organic Chemistry, 1961, (26), 2353-2355) was added dropwise to ethanolamine (325.8ml) at 40°C, with stirring under nitrogen and cooling to maintain the reaction temperature. The mixture was stirred for 3h whilst being heated at approximately 50°C before being cooled to room temperature. The mixture was partitioned between ethyl acetate and saturated aqueous sodium bicarbonate solution. The phases were separated and the organic phase was washed with further saturated aqueous sodium bicarbonate solution (100ml). The combined aqueous phases were further extracted with a portion of ethyl acetate (100ml) and the combined organic phases were washed with water (2x500ml) and saturated aqueous sodium chloride solution. The organic phase was concentrated *in vacuo* and re-concentrated after addition of toluene to give the title compound as a white solid (92.5g).
Mass Spectrum m/z 188 $[MH^+]$

Description 2: [4-(3,4-Difluorobenzyl)morpholin-2-yl]methylamine)




20

- 2-(Oxiran-2-ylmethyl)-1H-indole-1,3(2H)-dione (166.8g) was added portion-wise to the stirred compound of Description 1 (128.12g) at 80-90°C under nitrogen over 50mins. The mixture was stirred at 80-90°C for a further 3h before concentrated sulphuric acid (200ml) was added dropwise over 40mins. The mixture was heated to 150°C and stirred overnight. After cooling to room temperature, the mixture was washed with ethyl acetate (2x400ml). The aqueous phase was cooled to 10°C and saturated aqueous sodium chloride (400ml) was added carefully over 30mins. Further saturated aqueous sodium chloride (400ml) was added to the aqueous phase and the mixture was washed

PG4791


with further portions of ethyl acetate (3x500ml) The aqueous phase was cooled to 10°C and adjusted to pH 12-13 with 10N aqueous sodium hydroxide. The mixture was extracted with ethyl acetate (3x500ml) and the combined organic phases were filtered through 'Hyflo' filter aid. The combined organic extracts were washed with water (3x500ml) and saturated aqueous sodium chloride. The organic phase was concentrated *in vacuo* and re-concentrated (x3) after addition of toluene (3x150ml) to give the title compound as an amber oil (94g).
Mass Spectrum *m/z* 243 [MH⁺]

10 Description 3: 1-[(2S)-4-(3,4-Difluorobenzyl)morpholin-2-yl]methanamine



Description 2 (4.5g) (racemic mixture) was separated into its single enantiomers by preparative chiral-HPLC. The separation was carried out using a 2" x 22cm Chiralpak AD 20µm column, Merck self pack DAC system, eluting with 95:5:0.1 (v/v) heptane : absolute ethanol : diethylamine (flow rate: 50ml/min over 40min, UV detection 220nm); sample load preparation: 300mg sample in 10ml 1:1 (v/v) absolute ethanol : system eluent. The solvent was removed from the required fractions *in vacuo* and the product was partitioned between dichloromethane and 2N aqueous hydrochloric acid, the phases separated and the organic phase extracted with further 2N aqueous hydrochloric acid (x2). The combined aqueous extracts were adjusted to pH12 by the addition of solid potassium carbonate and the solution was saturated with solid sodium chloride. The solution was extracted with dichloromethane (x3) and the combined organic extracts washed with water, dried over magnesium sulphate, filtered and the solvent removed *in vacuo* to give the title compound (1.84g) as a yellow oil. Preparative HPLC retention time 28.5mins

30 Description 4: 3-[(methylsulphonyl)amino]phenylacetic acid



PG4791

To a stirred solution of 3-aminophenylacetic acid (3.2g) and sodium carbonate (5.44g) in de-ionised water (36ml) was added methanesulphonyl chloride (1.7 ml). The mixture was heated with stirring at 85° for 4h before being left to cool to room temperature, acidified to pH 2 with conc. hydrochloric acid and left to stand in a refrigerator at 4° overnight. The precipitated solid was filtered, washed with water and ether and the combined filtrate and washings were evaporated to dryness *in vacuo*. The resultant residue was dissolved in hot water and left to re-crystallise overnight whilst standing in a refrigerator at 4°C. The crystals were filtered, washed with a small quantity of cold water and dried *in vacuo* to give the title compound (0.417g) as colourless crystals.

LC/MS : $R_t = 2.00\text{min}$, m/z 228 [MH], m/z 247 [MNH₄⁺]


15

PG4791

Claims

1. A compound of formula (I):


5



and salts and solvates thereof.


2. A process for the preparation of a compound of formula (I) or a salt or solvate thereof which process comprises coupling of a compound of formula (II) or a salt thereof with a compound of formula (III)

10



(II)

15




(III)

3. A compound as claimed in claim (I) for use as a therapeutic agent


PG4791

4. A method of treatment of a human or animal subject suffering from or susceptible to an inflammatory condition which method comprises administering an effective amount of a compound as claimed in claim 1.
5. A pharmaceutical composition comprising a compound as claimed in claim 1 and a physiologically acceptable carrier therefor.
6. A compound of formula (II)



10

7. A compound of formula (IIBR)




15

wherein A is a protected amino group.


8. A compound of formula (IIB)

PG4791



wherein A is a protected amino group.

5 9. A compound of formula (V)



INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/03339

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/5375 C07D265/30

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00 35877 A (DU PONT PHARM CO (US)) 22 June 2000 (2000-06-22) page 6, line 26 - line 29 examples 58,125-133 ---	1,4,5
A	EP 0 243 959 A (DAINIPPON PHARMACEUTICAL CO LTD (JP)) 4 November 1987 (1987-11-04) claims ---	1-3,5-9
A	US 5 405 854 A (SCHMID GERARD ET AL) 11 April 1995 (1995-04-11) column 7, line 54 - line 65 example 7B ---	2,6
P,X	WO 02 26722 A (GLAXO GROUP LTD (GB)) 4 April 2002 (2002-04-04) page 3, line 28; example 96 ---	1-6
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* & * document member of the same patent family

Date of the actual completion of the international search

2 July 2003

Date of mailing of the international search report

14/07/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 551 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Cortés, J

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 03/03339

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 02 26723 A (GLAXO GROUP LTD (GB)) 4 April 2002 (2002-04-04) intermediate 21, page 35 -----	6

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/03339

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0035877	A	22-06-2000	AU 2056800 A CA 2347912 A1 EP 1140834 A1 WO 0035877 A1 US 6331545 B1 US 2002119980 A1	03-07-2000 22-06-2000 10-10-2001 22-06-2000 18-12-2001 29-08-2002
EP 0243959	A	04-11-1987	AT 79616 T AU 592177 B2 AU 7227587 A CA 1291135 C CZ 8702897 A3 DD 256325 A5 DE 3781195 D1 DE 3781195 T2 DK 218787 A EP 0243959 A1 ES 2043617 T3 FI 871828 A ,B, GR 3005934 T3 HK 18194 A HU 45514 A2 HU 198193 B JP 1686407 C JP 3054937 B JP 63264467 A JP 2042069 A MX 6263 A NZ 220120 A PH 24049 A SG 494 G SK 289787 A3 SU 1597101 A3 US 4870074 A ZA 8703022 A KR 9508530 B1	15-09-1992 04-01-1990 05-11-1987 22-10-1991 11-11-1998 04-05-1988 24-09-1992 18-02-1993 31-10-1987 04-11-1987 01-01-1994 31-10-1987 07-06-1993 11-03-1994 28-07-1988 28-08-1989 11-08-1992 21-08-1991 01-11-1988 13-02-1990 01-11-1993 28-06-1989 05-03-1990 15-04-1994 03-06-1998 30-09-1990 26-09-1989 20-10-1987 31-07-1995
US 5405854	A	11-04-1995	AT 203013 T AU 3391693 A BG 61506 B1 BG 97493 A BR 9300753 A CA 2089972 A1 CN 1076690 A CZ 9300346 A3 DE 59310186 D1 DK 559046 T3 EP 0559046 A1 ES 2161217 T3 FI 930990 A FI 991361 A GR 3036870 T3 HR 930266 A1 HU 70156 A2 HU 9500254 A3 IL 120727 A JP 2045220 C JP 6025195 A	15-07-2001 09-09-1993 31-10-1997 24-03-1994 08-09-1993 07-09-1993 29-09-1993 15-12-1993 16-08-2001 05-11-2001 08-09-1993 01-12-2001 07-09-1993 14-06-1999 31-01-2002 30-04-1996 28-09-1995 28-09-1995 16-08-1998 09-04-1996 01-02-1994

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/03339

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5405854	A		JP 7080848 B	30-08-1995
			KR 274422 B1	15-12-2000
			MX 9301155 A1	01-09-1993
			NO 930819 A	07-09-1993
			NZ 247014 A	27-11-1995
			PL 297960 A1	20-09-1993
			PT 559046 T	30-01-2002
			RO 112863 B	30-01-1998
			RU 2133739 C1	27-07-1999
			SG 48063 A1	17-04-1998
			SI 9300104 A	30-09-1993
			SK 14893 A3	10-11-1993
			US 5578594 A	26-11-1996
			US 5677448 A	14-10-1997
			US 5763604 A	09-06-1998
			ZA 9301403 A	06-09-1993
WO 0226722	A	04-04-2002	AU 9014301 A	08-04-2002
			WO 0226722 A1	04-04-2002
WO 0226723	A	04-04-2002	AU 9014601 A	08-04-2002
			WO 0226723 A1	04-04-2002